arXiv:1210.4150v1 [math.PR] 15 Oct 2012

New methods to determine the critical probability in fractal percolation

Henk Don

October 16, 2012

1 Introduction

Fractal percolation has been introduced by Mandelbrot in 1974 as a model for turbulence and is discussed in his book *The Fractal Geometry of Nature* [7]. Several equivalent formal definitions of this process can be found in the literature (see e.g. [2, 3, 5]). Here we only give an informal definition of the two-dimensional case. Let K_0 be the unit square and choose an integer $M \geq 2$ and a parameter $p \in [0,1]$. To obtain K_1, divide K_0 into M^2 equal subsquares, each of which survives with probability p and is discarded with probability $1-p$, independently of all other subsquares. Now do the same procedure in all surviving squares, in order to obtain K_2. Iterating this process gives a decreasing sequence of sets $(K_n)_{n\in\mathbb{N}}$, see Figure 1. Let $K = \bigcap_{n\in\mathbb{N}} K_n$ be the limit set.



Figure 1: Realizations of $K_n, n = 1, 2, 3, 5, 7$ for $M = 3$ and $p = 0.85$.

It was shown in 1988 by Chayes, Chayes and Durrett [2] that there exists a non-trivial critical value $p_c(M)$ such that a.s. the largest connected component in K is a point for $p < p_c(M)$ and with positive probability there is a connected component intersecting opposite sides of the unit square for $p \geq p_c(M)$.

For all $M \geq 2$, the value of $p_c(M)$ is unknown. Several attempts have been made to find bounds for $p_c(M)$. It is easy to see that K is empty a.s. if $p \leq 1/M^2$, which implies $p_c(M) > 1/M^2$. The argument in [2] is already a bit smarter: any left-right crossing has to cross the line $\{1/M\} \times [0,1]$ somewhere. A crossing of this line in K_n means that there is a pair of adjacent squares on opposite sides of this line. Such pairs form a branching process with mean offspring p^2M and consequently $p_c(M) > 1/\sqrt{M}$. For the case $M = 2$ this was sharpened by White in 2001 to $p_c(2) \geq 0.810$, who used a set that dominates K and has a simpler structure to study.

Sharp upper bounds are harder to obtain. The first idea to get rigorous upper bounds for $M \geq 2$ was given by Chayes, Chayes and Durrett [2], but (in their own words) these bounds are ridiculously close to 1. For $M = 3$, they show that $p_c(3) < 0.9999$ (although in fact one can prove that $p_c(3) < 0.993$ with their method), which was improved by Dekking and Meester [4] to $p_c(3) < 0.991$. Chayes et al only treat $M = 3$, but they point out that the same idea works for any $M \geq 3$. The case $M = 2$ can be treated by comparing with $M = 4$. As is noted by van der Wal [9], a coupling argument gives $p_c(2) \leq 1-(1-\sqrt{p_c(4)}\,)^4$. Following this approach gives $p_c(4) < 0.998$ and $p_c(2) < 1 - 10^{-12}$.

In this paper we present ideas to find significantly sharper lower and upper bounds. To find lower bounds, we compare fractal percolation with site percolation. A fundamentally new result is that for all M we construct an increasing sequence that converges to $p_c(M)$. The terms in the sequence can in principle be calculated algorithmically. To actually obtain numerical results we bound them from below. These ideas lead to (computer aided) proofs that $p_c(2) > 0.881$ and $p_c(3) > 0.784$.

For the upper bounds, we map all possible realizations of K_n to a finite alphabet $\mathscr{A}$. The choice of the alphabet $\mathscr{A}$ is inspired by van der Wal's work [9]. The fractal percolation iteration process now induces an iterative random process on $\mathscr{A}$, which is easier to analyze than the original process. This theoretical framework is the basis of computer aided proofs for the following upper bounds: $p_c(2) < 0.993$, $p_c(3) < 0.940$ and $p_c(4) < 0.972$.

2 Lower bounds for $p_c(M)$

In this section we develop methods to calculate lower bounds for the critical value of two-dimensional fractal percolation. First we briefly introduce site percolation and then we prove a coupling with fractal percolation that allows us to find lower bounds for $p_c(M)$. In particular, we construct an increasing sequence of lower bounds and we prove that this sequence converges to $p_c(M)$. At the end of this section we show how to use these insights to obtain numerical results.

2.1 Site percolation

Consider the infinite two-dimensional square lattice in which each vertex is open with probability p and closed otherwise. In this model the percolation probability $\zeta(p)$ is defined as the probability that the origin belongs to an infinite open cluster. The critical probability is given by

$$p_c^{site} := \inf \{p : \zeta(p) > 0\}.$$

It has been shown by van den Berg and Ermakov [1] that $p_c^{site} > 0.556$. Now consider the probability that a box of $n \times n$ vertices is crossed by an open cluster. This probability is monotone in n, so it converges to a limit. It is a classical result that a positive percolation probability is equivalent to this limit being positive (see e.g. [6] and the references therein). As a consequence we have the following property, that will be used to couple site percolation to fractal percolation.

Property 1 *Take a box of $n \times n$ vertices. Suppose $p < p_c^{site}$. Then the probability that there is an open cluster intersecting opposite sides of the box converges to 0 as $n \to \infty$.*

2.2 Coupling site percolation and fractal percolation

In fractal percolation, a set in the unit square is said to percolate if it contains a connected component intersecting opposite sides of the square. Let

$$\theta_n(p, M) = \mathbb{P}(K_n(p, M) \text{ percolates}), \qquad \theta(p, M) = \mathbb{P}(K(p, M) \text{ percolates}).$$

Then $p_c(M) := \inf \{p : \theta(p, M) > 0\}$. We will often suppress some of the dependence on M and p. It is well known that $\lim_{n\to\infty} \theta_n(p) = \theta(p) = \mathbb{P}(\bigcap_{n=0}^{\infty} \{K_n(p) \text{ percolates}\})$, see [8].
To obtain a proper coupling, we have to deal with diagonal connections. For example, the set $[0, 1/2]^2 \cup [1/2, 1]^2$ percolates. We would like to ignore such diagonal connections, since in site percolation diagonal connections do not exist. Therefore we redefine percolation as follows. We say K_n percolates if there is a left-right crossing of the square that does not use diagonal connections. Since diagonal connections break down almost surely, this does not change the limiting percolation probability. Also from now on connections, crossings and connected components in K_n are similarly redefined.

Theorem 1 *Let $\pi_n(p, M) = \mathbb{P}($at least two sides are connected in $K_n(p, M))$. If $\pi_n(p) < p_c^{site}$ for some n, then $p < p_c(M)$.*

Proof First define *delayed* fractal percolation: $F_{m,n}$ is constructed in the same way as K_{m+n}, the only difference being that we do not discard any squares in the first m construction steps. So we first divide the unit square into $M^m \times M^m$ subsquares and only then we start the fractal percolation process in each of these squares. Delayed fractal percolation stochastically dominates fractal percolation.



Figure 2: Three coupled realizations to illustrate (1). Left: K_5 for $M = 2$ and $p = 2/3$. Middle: $F_{2,3}$ for $M = 2$ and $p = 2/3$. Right: K_1 for $M = 4$ and $p = \pi_3(2/3, 2)$.

Then we have the following inequalities (illustrated for $M = m = 2$ and $n = 3$ in Figure 2):

$$\theta_{m+n}(p, M) \leq \mathbb{P}(F_{m,n}(p, M) \text{ percolates}) \leq \theta_1(\pi_n(p), M^m). \tag{1}$$

The first inequality follows from the fact that $F_{m,n}$ is probabilistically larger than K_{m+n}. The second inequality can be explained as follows. Suppose we have two types of squares: realizations of $K_n(p, M)$ in which at least two sides are connected (type 1) and realizations in which no side is connected to any other side (type 2). Suppose we tile a larger square with M^{2m} independent realizations of $K_n(p, M)$. So the probability on a type 1 square is $\pi_n(p)$. This larger square is a (scaled) realization of $F_{m,n}(p, M)$. Now replace all type 1 squares by a full square and discard all type 2 squares. This gives a realization of $K_1(\pi(p), M^m)$. Moreover, this replacement procedure can not destroy percolation. To see this, suppose we have a left-right crossing in $F_{m,n}(p, M)$. All parts of this crossing that are in type 1 squares are preserved. If the crossing uses a type 2 square S, then it enters and leaves S at the same side of S. At this side the neighboring square T must be of type 1. So if we remove S and replace T by a full square, there still is a left-right crossing.

A first level fractal percolation set can be seen as site percolation in a finite box. Suppose $\pi_n(p) < p_c^{site}$ for some n and let the box size M^m tend to ∞. By Property 1 we arrive at

$$\lim_{m\to\infty} \theta_1(\pi_n(p), M^m) = 0. \tag{2}$$

Therefore, if $\pi_n(p) < p_c^{site}$, by (1) we find that

$$\theta(p) = \lim_{m\to\infty} \theta_{m+n}(p) = 0,$$

which in turn implies $p < p_c(M)$. □

2.3 A convergent sequence of lower bounds for $p_c(M)$

In this section we define a sequence of lower bounds for $p_c(M)$. We prove that this sequence converges to the $p_c(M)$. Let

$$p_c^n(M) = \sup\left\{p : \pi_n(p, M) < p_c^{site}\right\}. \tag{3}$$

Note that $\pi_n(p_c^n) \leq p_c^{site}$ for all n, since $\pi_n(p)$ is continuous in p. Theorem 1 requires a strict inequality, but this does not give any problem since $\pi_n(p)$ is strictly decreasing in n, so $\pi_{n+1}(p_c^n) < p_c^{site}$ for all n and hence indeed $p_c^n(M) < p_c(M)$ for all n. The strict monotonicity in n also implies that $(p_c^n(M))_{n=0}^{\infty}$ is increasing. The obvious question now is whether $(p_c^n(M))_{n=0}^{\infty}$ converges to $p_c(M)$. We will show that this is indeed the case. First we need that $\pi_n(p)$ goes to zero if the fractal percolation is subcritical:

Lemma 1 *If $p < p_c(M)$, then $\lim_{n\to\infty} \pi_n(p, M) = 0$.*

Proof Suppose $p < p_c(M)$, so $\theta(p) = 0$. Note that a.s. there is an n such that in K_n the two squares in the top left and bottom left corner are discarded already. Conditioned on this event, a connection in the limiting set from the left side to any other side can only occur if it horizontally crosses the vertical strip S consisting of the squares $[0, M^{-n}] \times [jM^{-n}, (j+1)M^{-1}]$ for $j = 1, \ldots, M^n - 2$. From selfsimilarity and subcriticality it follows that in the limit each of these squares has zero probability to contain a component connecting opposite sides (horizontally and vertically). So in K a horizontal crossing of S can only occur if it crosses a block of two vertically adjacent squares. But as Dekking and Meester showed (Lemma 5.1 in [4]), $\theta(p) = 0$ implies that such a block crossing has zero probability as well. It follows that connections from the left side to any other side have zero probability to occur. If this holds for the left side, then it holds for all sides, and so $\lim_{n\to\infty} \pi_n(p) = 0$. □

The previous lemma makes it easy to prove convergence:

Theorem 2 *The sequence $(p_c^n(M))_{n=0}^{\infty}$ converges to $p_c(M)$ if $n \to \infty$.*

Proof Let $\varepsilon > 0$ and suppose $p = p_c(M) - \varepsilon$. Then by Lemma 1, there exists an N such that $\pi_n(p) < p_c^{site}$ for all $n \geq N$. Therefore, $p_c^n(M) > p$ for all $n \geq N$. □

The theory developed so far gives us an algorithmic tool to calculate an increasing and converging sequence of lower bounds for $p_c(M)$. Still, it is not so easy to actually obtain sharp bounds. For example, a little thought gives

$$\begin{aligned}\pi_1(p,2) &= 1-(1-p)^4,\\ \pi_1(p,3) &= 1+(1-p)^4(p^5+4p^4(1-p)+6p^3(1-p)^2-1),\end{aligned}$$

so using the bound of van den Berg and Ermakov for p_c^{site}, we find $p_c^1(2) > 0.183$ and $p_c^1(3) > 0.178$. Of course, we want to find a bit sharper bounds, so we should take larger values of n. However, for large n, the functions $\pi_n(p)$ are very complicated polynomials, and it is not clear how to find them in reasonable time. In the next section we will discuss a way to avoid this problem.

3 Classifying realizations

In this section we introduce the idea to map realizations of K_n to a finite alphabet $\mathscr{A}$ that does not depend on n. Such a map will be called a *classification*. In this way, we can simplify the substitution process without losing too much essential information on the connectivity structure in K_n. The construction of the random sets $(K_n)_{n=0}^\infty$ now induces a sequence of probability measures on $\mathscr{A}$. These ideas can be used to obtain both lower and upper bounds for $p_c(M)$. For the lower bounds we choose the classification in such a way that we can calculate upper estimates for $\pi_n(p)$ and then we can approximate terms of $(p_c^n)_{n=0}^\infty$ using the van den Berg and Ermakov bound for p_c^{site}. For the upper bounds we will construct a classification that permits us to bound $\theta(p)$ away from 0 for p sufficiently large.
Let $\mathscr{K}_n$ be the set of all possible realizations of K_n. We will define a partially ordered alphabet $\mathscr{A}$ and the classification will actually be a sequence of maps $\mathscr{C} = (\mathscr{C}_n)_{n=0}^\infty$, where $\mathscr{C}_n$ is a map from $\mathscr{K}_n$ to $\mathscr{A}$. The alphabet will have a unique minimum and maximum, with the property that

$$\mathscr{C}_n(\emptyset) = \min(\mathscr{A}) \quad \text{and} \quad \mathscr{C}_n([0,1]^2) = \max(\mathscr{A}). \tag{4}$$

Realizations of K_n can be mapped in a natural way to words in $\mathscr{A}^{M\times M}$. A realization of K_n is obtained by first generating a realization of K_1 and then replacing all squares that survived by independent scaled realizations of K_{n-1}. Given $\mathscr{C}_{n-1}$, this induces a map $\mathscr{W}_n$ from $\mathscr{K}_n$ to $\mathscr{A}^{M\times M}$. If $\mathscr{C}_n(K_n)$ only depends on the word $\mathscr{W}_n(K_n)$ and not on n, we say the classification is *regular*:

Definition 1 *Let ϕ be a map from $\mathscr{A}^{M\times M}$ to $\mathscr{A}$. Define $\mathscr{C} = (\mathscr{C}_n)_{n=0}^\infty$ by letting*

$$\mathscr{C}_n = \phi \circ \mathscr{W}_n, \qquad n \geq 1. \tag{5}$$

Then we say $\mathscr{C}$ is a regular classification *and ϕ is called the* word code *of $\mathscr{C}$.*

Example 1 Let $M = 2$ and take the alphabet $\mathscr{A} = \{\square, \boxplus\}$, where $\min \mathscr{A} = \square$ and $\max \mathscr{A} = \boxplus$. For a 2×2 word w over $\mathscr{A}$ let $\phi(w) = \boxplus$ if and only if at least one of the letters in w equals $\boxplus$. Let $\mathscr{C} = (\mathscr{C}_n)_{n=0}^\infty$ be regular with word code ϕ. Then $\mathscr{C}_0$ is determined by (4):

$$\mathscr{C}_0(K_0(p)) = \mathscr{C}_0([0,1]^2) = \boxplus.$$

For $n \geq 1, \mathscr{C}_n = \phi \circ \mathscr{W}_n$. For instance

$$\mathscr{C}_1([\frac{1}{2},1]^2) = \phi(\mathscr{W}_1([\frac{1}{2},1]^2)) = \phi \left\{ \begin{array}{cc} \mathscr{C}_0(\emptyset) & \mathscr{C}_0([0,1]^2) \\ \mathscr{C}_0(\emptyset) & \mathscr{C}_0(\emptyset) \end{array} \right\} = \phi \left\{ \begin{array}{cc} \square & \boxplus \\ \square & \square \end{array} \right\} = \boxplus.$$

$\blacksquare$

We now want to analyze the probabilities $\mathbb{P}(\mathscr{C}(K_n(p)) = a)$, where $a \in \mathscr{A}$. Suppose $\mathscr{C}$ is a regular classification with word code ϕ. Let $x \in [0,1]^{|\mathscr{A}|}$ be a probability vector on $\mathscr{A}$, and suppose we construct an $M \times M$ word w in which all letters are chosen independently according to x. Define $F_{\mathscr{C}}(x) \in [0,1]^{|\mathscr{A}|}$ by

$$(F_{\mathscr{C}}(x))_a = \mathbb{P}_x(\phi(w) = a), \qquad a \in \mathscr{A}.$$

The function $F_{\mathscr{C}} : [0,1]^{|\mathscr{A}|} \to [0,1]^{|\mathscr{A}|}$ will be the key to calculate the probabilities $\mathbb{P}(\mathscr{C}(K_n(p)) = a)$ in an iterative way, as is shown in the next lemma. Let $\tau^n(p)$ be a vector indexed by the elements of $\mathscr{A}$ such that

$$\tau^n_a(p) = \mathbb{P}(\mathscr{C}(K_n(p)) = a).$$

Let $\tau^{\square}$ and $\tau^{\boxplus}$ be the vectors that assign full probability to $\min(\mathscr{A})$ and $\max(\mathscr{A})$ respectively.

Lemma 2 *If the classification $\mathscr{C}$ is regular, then*

$$\tau^{n+1}(p) = F_{\mathscr{C}}(p\tau^n(p) + (1-p)\tau^{\square}) \quad \textit{with initial condition} \quad \tau^0(p) = \tau^{\boxplus}.$$

Proof The letters in $\mathscr{W}_{n+1}(K_{n+1}(p))$ are independent. With probability p a letter corresponds to a scaled realization of $K_n(p)$, with probability $1-p$ it corresponds to an empty square. So each letter occurs according to the probability vector $p\tau^n(p) + (1-p)\tau^{\square}$. This gives the recursion. The initial condition follows from (4). $\square$

This recursion formula is essentially a generalization of the recursion given in [4].

3.1 Strategy for lower bounds

The strategy to find lower bounds for p_c now is as follows. Define an alphabet $\mathscr{A}$ with subset $\mathscr{A}_\pi$ and define a classification $\mathscr{C}$ (by choosing ϕ) in such a way that $\mathbb{P}(\mathscr{C}(K_n(p)) \in \mathscr{A}_\pi) \geq \pi_n(p)$ for all n. Now search for the largest p for which this probability is smaller than 0.556. We will give an example to illustrate this procedure. In this example (which only gives a very moderate bound) $\mathscr{A}$ only has two elements, and therefore exactly the same recursion as in [4] pops up.

Example 2 (A lower bound for $M = 2$) Let $\mathscr{C}$ be the classification of Example 1. By induction it follows that if two sides are connected in K_n, then $\mathscr{C}_n(K_n) = \boxplus$. Consequently $\tau_{\boxplus}^{n}(p) \geq \pi_n(p)$. From the definition of ϕ it follows that in this case

$$\tau_{\boxplus}^{n+1}(p) = 1 - (1 - p\tau_{\boxplus}^{n}(p))^4 = \tau_{\boxplus}^{1}(p\tau_{\boxplus}^{n}(p)).$$

The function $G_p(x) := \tau_{\boxplus}^{1}(px)$ is increasing and $\tau_{\boxplus}^{n}(p)$ decreases to the largest fixed point of $G_p(\cdot)$. Choosing $p = 0.33$, we find $\pi_{50}(p) \leq \tau_{\boxplus}^{50}(p) \approx 0.554 < 0.556 \leq p_c^{site}$, and consequently $p_c^{50}(2) > 0.33$ and therefore also $p_c(2) > 0.33$. $\blacksquare$

3.2 Strategy for upper bounds

Our recipe to find upper bounds for $p_c(M)$ is a bit more involved. We already have a partial ordering on $\mathscr{A}$, but to find upper bounds it will also be convenient to partially order the probability vectors on $\mathscr{A}$. A set $S \subseteq \mathscr{A}$ will be called *increasing* if $a \in S$ implies $b \in S$ for all $b \succeq a$. For probability vectors x and y, we now write $x \succeq y$ if x assigns larger probabilities to all increasing subsets of $\mathscr{A}$. We say the function $F_{\mathscr{C}}$ is increasing if $F_{\mathscr{C}}(x) \succeq F_{\mathscr{C}}(y)$ for $x \succeq y$.

Lemma 3 *Let $\mathscr{C}$ be a regular classification for which $F_{\mathscr{C}}$ is increasing. Then $(\tau^n(p))_{n=0}^{\infty}$ is decreasing and $\tau^{\infty}(p) := \lim_{n\to\infty} \tau^n(p)$ exists. If $F_{\mathscr{C}}(px + (1-p)\tau^{\square}) \succeq x$ for some probability vector $x \in [0,1]^{|\mathscr{A}|}$ and $p \in (0,1]$, then $\tau^{\infty}(p) \succeq x$.*

Proof Since $\tau^0(p) = \tau^{\boxplus}$, we have $\tau^0(p) \succeq \tau^1(p)$. Suppose $\tau^n(p) \succeq \tau^{n+1}(p)$. Then $\tau^{n+1}(p) = F_{\mathscr{C}}(p\tau^n(p) + (1-p)\tau^{\square}) \succeq F_{\mathscr{C}}(p\tau^{n+1}(p) + (1-p)\tau^{\square}) = \tau^{n+2}(p)$ since $F_{\mathscr{C}}$ is increasing. So $(\tau^n(p))_{n=0}^{\infty}$ is decreasing. If $S \subseteq \mathscr{A}$ is an increasing set, then

$$\tau_S^n(p) := \mathbb{P}(\mathscr{C}(K_n) \in S)$$

decreases in n and is bounded from below by 0, so $\lim_{n\to\infty} \tau_S^n(p)$ exists. There are only finitely many increasing subsets of $\mathscr{A}$ and their limiting probabilities uniquely determine $\tau^{\infty}(p)$.
For the second statement, $\tau^0(p) = \tau^{\boxplus} \succeq x$. Now suppose $\tau^n(p) \succeq x$ for some n. Then

$$\tau^{n+1}(p) = F_{\mathscr{C}}(p\tau^n(p) + (1-p)\tau^{\square}) \succeq F_{\mathscr{C}}(px + (1-p)\tau^{\square}) \succeq x.$$

and therefore $\tau^n(p) \succeq x$ for all $n \in \mathbb{N}$. Hence $\tau^{\infty}(p) \succeq x$. $\square$

Corollary 1 *Let $\mathscr{C}$ be a regular classification for which $F_{\mathscr{C}}$ is increasing. Suppose $x \in [0,1]^{|\mathscr{A}|}$ is a probability vector for which $x_{\max(\mathscr{A})} > 0$. If $\theta_n(p) \geq \tau_{\max(\mathscr{A})}^n(p)$ for all n, and $F_{\mathscr{C}}(px + (1-p)\tau^{\square}) \succeq x$ for some $p \in (0,1]$, then*

$$\theta(p) = \lim_{n\to\infty} \theta_n(p) \geq \lim_{n\to\infty} \tau_{\max(\mathscr{A})}^n(p) \geq x_{\max(\mathscr{A})} > 0,$$

and consequently $p > p_c$.

Before we give an example of the procedure to find upper bounds for $p_c(M)$, we will first construct suitable alphabets in the next section.

4 Construction of the alphabet and word codes

Our alphabets will be defined by means of non-crossing equivalence relations. Figure 3 displays all equivalence relations on an ordered set of 5 elements, of which 42 are non-crossing. Let $E = \{e_1, \ldots, e_n\}$ be a collection of line segments that together form the boundary of the unit square, numbered clockwise, starting from $(0,0)$. We define the alphabet $\mathscr{A}_E$ as the set of all non-crossing equivalence relations on E. The size of $\mathscr{A}_E$ is given by the Catalan number $\frac{1}{n+1}\binom{2n}{n}$.
On the alphabet $\mathscr{A}_E$, we can naturally define a partial ordering. For $a, b \in \mathscr{A}_E$ we say $a \succeq b$ if all equivalences that hold in b also hold in a. One should think of the equivalences (denoted by $\sim$) as connections between the elements. The letters can be represented as pictures in a square. For example, if E is the set containing the four sides of the square (so $n = 4$), then $\mathscr{A}_E$ can be represented as



Figure 3: Equivalence relations on 5 elements

$\mathscr{A}_E =$ {□, ◸, ◹, ◿, ◺, ◫, ⊟, ⧄, ⧅, ⊤, ⊢, ⊥, ⊣, ⊞} .

The alphabet has a unique maximum and minimum, $\max(\mathscr{A}_E) =$ ⊞ and $\min(\mathscr{A}_E) =$ □. From now on we will denote the alphabet that corresponds to dividing each side in M^n elements of equal size by $\mathscr{A}_{M,n}$.
Now that we have letters, we can start making words. We only allow square words, obtained by tiling $[0, M]^2$ by M^2 letters. An example of a 3×3 word over $\mathscr{A}_{3,0}$ is given in the right panel of Figure 4. A word w also has a boundary element set E_w. We enumerate clockwise, starting from $(0,0)$, as follows: $E_w = \left\{e_1^w, \ldots, e_{M|E|}^w\right\}$.

4.1 Weak and strong connectivity

Suppose we partition E_w into subsets $E_w^1, \ldots, E_w^k$ for some integer $k \geq 2$. If $e \sim f$ for some $e \in E_w^i$ and $f \in E_w^j$, we say E_w^i and E_w^j are *connected*. If $E_w^{i_1}, \ldots, E_w^{i_m}$ form a chain of pairwise connected sets, we say $E_w^{i_1}$ and $E_w^{i_m}$



Figure 4: Left: A realization of K_2 for $M = 3$. Right: A 3×3 word w over $\mathscr{A}_{3,0}$. If we use the word code $\Psi_{3,0}$, then this word corresponds to the realization on the left.

are *weakly connected*. If strictly more than half of the elements in E_w^i and strictly more than half of E_w^j are in the same equivalence class, we say E_w^i and E_w^j are *strongly connected*. Observe that weak and strong connectivity are equivalence relations on $\{E_w^1, \ldots, E_w^k\}$.
Choose the alphabet $\mathscr{A}_{M,n}$, then $|E| = 4M^n$, so words of size $M \times M$ have $4M^{n+1}$ boundary elements. Partition E_w in the subsets

$$E_w^i = \left\{ e^w_{M(i-1)+j} : 1 \leq j \leq M \right\}, \quad 1 \leq i \leq 4M^n.$$

This gives a tool to define word codes: a word defines an equivalence relation on $\{E_w^1, \ldots, E_w^{4M^n}\}$ that obviously can be mapped to an equivalence relation on $E = \{e_1, \ldots, e_{4M^n}\}$, which is just a letter in $\mathscr{A}_E$. In this way, define word codes

$$\Phi_{M,n} : \mathscr{A}_{M,n}^{M \times M} \to \mathscr{A}_{M,n} \quad \text{and} \quad \Psi_{M,n} : \mathscr{A}_{M,n}^{M \times M} \to \mathscr{A}_{M,n}$$

based on weak and strong connectivity respectively. The corresponding classifications are increasing.

Example 3 Let w be the word in Figure 4. This word has 12 boundary elements that will be grouped into four partition sets. For instance, E_w^3 contains the three elements at the right side. Then E_w^1, E_w^2 and E_w^4 are all pairwise connected and E_w^3 is connected to E_w^4. Consequently E_w^i is weakly connected to E_w^j for all i and j. Therefore $\Phi_{3,0}(w) = \boxplus$.
The only partition sets that are strongly connected are E_w^1 and E_w^2. Therefore $\Psi_{3,0}(w) =$ ◰. ■

The idea behind the definitions of $\Phi_{M,n}$ and $\Psi_{M,n}$ is that they guarantee the following key properties:

Property 2 *Using the alphabet $\mathscr{A}_{M,k}$ and word code $\Phi_{M,k}$, the following implication holds: if $e_i, e_j \in E$ are connected in K_n, then $e_i \sim e_j$ in $\mathscr{C}(K_n)$.*

Property 3 *Using the alphabet $\mathscr{A}_{M,k}$ and word code $\Psi_{M,k}$, the reversed implication holds: if $e_i \sim e_j$ in $\mathscr{C}(K_n)$, then e_i and e_j are connected in K_n.*

These properties are not hard to prove. Therefore we do not give a fully detailed proof. For Property 2 a straightforward inductive argument suffices. For Property 3, one should note that we actually have a stronger implication, namely that e_i and e_j are connected in a special way that we will explain using the realization in Figure 4. In this realization the left side and the top side are connected. At both these sides there are two (strictly more than $M/2$) first level squares each of which contains two second level squares that survived. These second level squares are all in the same connected component. This property is easily generalized to higher level realizations and other values of M. If two neighbouring squares both have such connection from their joint side to an other side, then these two connections are in the same connected component.

The following lemma shows that these alphabets and word codes are suitable for our purposes, see the discussion in Section 3.

Lemma 4 *Using the word codes $\Phi_{M,k}$ and $\Psi_{M,k}$ leads to two inequalities:*

1. *Take the alphabet $\mathscr{A}_{M,k}$ and define a classification $\mathscr{C}$ by the word code $\Phi_{M,k}$. Let $\mathscr{A}_\pi \subseteq \mathscr{A}_{M,k}$ be the set of all letters in which at least two of the sides are connected. Then $\tau_\pi^n(p) := \mathbb{P}(\mathscr{C}(K_n(p)) \in \mathscr{A}_\pi) \geq \pi_n(p)$ for all n.*
2. *Take the alphabet $\mathscr{A}_{M,k}$ and define a classification $\mathscr{C}$ by the word code $\Psi_{M,k}$. Then $\tau^n_{\max(\mathscr{A})}(p) \leq \theta_n(p)$ for all n.*

Proof These statements follow from Property 2 and 3 respectively. $\square$

Now we are ready to give an example illustrating how to find upper bounds. We will keep the example as simple as possible, so that it can be checked by hand. Therefore our alphabet will contain only two letters and we will use a word code that is even simpler than $\Psi_{3,0}$. Nevertheless, it leads to a bound that already improves upon the best bound known so far.

Example 4 (An upper bound for $M = 3$) Let $\mathscr{A} = \{\max(\mathscr{A}_{3,0}), \min(\mathscr{A}_{3,0})\} = \{\square, \boxplus\}$. A 3×3 word w has twelve boundary elements that we partition into four sets as before. Define a classification $\mathscr{C}$ by letting $\phi(w) = \boxplus$ if and only if all sets $E_w^1, \dots, E_w^4$ are strongly connected. This classification is increasing and satisfies Property 3, so $\tau^n_{\max(\mathscr{A})}(p) \leq \theta_n(p)$ for all n. The recursion of Lemma 2 reduces to $\tau_{\boxplus}^{n+1}(p) = \tau_{\boxplus}^1(p\tau_{\boxplus}^n(p))$, where

$$\tau_{\boxplus}^1(p) = p^9 + 9p^8(1-p) + 20p^7(1-p)^2.$$

This function is the first component of $F_{\mathscr{C}}(p\tau^{\boxplus} + (1-p)\tau^{\square}) = F_{\mathscr{C}}((p, 1-p))$. Now choose $p = 0.984$ and $x = 0.9720$. Then $\tau_{\boxplus}^1(px) \approx 0.9721 > x$. Therefore the probability vector $(x, 1-x)$ satisfies the requirement of Corollary 1. Hence $p_c(3) < 0.984$. ■

4.2 Monotonicity and convergence

So far we developed some tools to find bounds for $p_c(M)$. One would expect that taking larger alphabets results in sharper bounds, since we can approximate the connectivity structure in K_n more accurately. In this section we show that this is indeed the case and that the lower bounds even convergence to $p_c(M)$. Unfortunately, we do not know if the upper bounds also converge to $p_c(M)$.
For the word code $\Phi_{M,k}$ over $\mathscr{A}_{M,k}$, define the corresponding classification and let $\tau_\pi^n(p)$ be defined as before. Then define a critical value as follows:

$$p_c(\Phi_{M,k}) := \sup\left\{p : \tau_\pi^\infty(p) < p_c^{site}\right\}.$$

Let $\mathscr{A} = \mathscr{A}_{M,k}$ and define $\mathscr{C}$ by the word code $\Psi_{M,k}$. Also here we define a critical value:

$$p_c(\Psi_{M,k}) := \inf\left\{p : \tau^\infty_{\max \mathscr{A}}(p) > 0\right\}.$$

Now we have the following proposition:

Proposition 1 *The sequence* $(p_c(\Phi_{M,k}))_{k=0}^\infty$ *is increasing and* $(p_c(\Psi_{M,k}))_{k=0}^\infty$ *is decreasing. Moreover,*

$$\lim_{k\to\infty} p_c(\Phi_{M,k}) = p_c(M).$$

Proof The alphabet $\mathscr{A}_{M,k}$ contains equivalence relations on an element set $E_{M,k}$. The element set $E_{M,k+1}$ of $\mathscr{A}_{M,k+1}$ can be obtained by dividing each element $e \in E_{M,k}$ into M equal pieces. These pieces will be called children of e, and e will be called the parent of its children. Define classifications $\mathscr{C}^k$ and $\mathscr{C}^{k+1}$ by the word codes $\Phi_{M,k}$ and $\Phi_{M,k+1}$. Denote the corresponding probability vectors by ${}^k\tau$ and ${}^{k+1}\tau$. By induction it follows that if two elements are connected in $\mathscr{C}^{k+1}(K_n)$, then their parents are connected in $\mathscr{C}^k(K_n)$. Consequently, ${}^k\tau_\pi^n(p) \geq {}^{k+1}\tau_\pi^n(p)$ and so

$$p_c(\Phi(M, k+1)) \geq p_c(\Phi(M, k)).$$

Now define $\mathscr{C}^k$ and $\mathscr{C}^{k+1}$ by the word codes $\Psi_{M,k}$ and $\Psi_{M,k+1}$. If two elements are connected in $\mathscr{C}^k(K_n)$, then two of their children are connected in $\mathscr{C}^{k+1}(K_n)$. Henceforth, ${}^k\tau^n_{\max \mathscr{A}}(p) \leq {}^{k+1}\tau^n_{\max \mathscr{A}}(p)$, so

$$p_c(\Psi(M, k+1)) \leq p_c(\Psi(M,k)).$$

Take the alphabet $\mathscr{A}_{M,k}$ and define $\mathscr{C}^k$ by the word code $\Phi_{M,k}$. A realization of K_n consists of $M^n \times M^n$ squares, and letters in $\mathscr{A}_{M,k}$ have M^k boundary elements at each side. This means that for $n \leq k$ the classification describes the connectivity structure exactly: two elements in $\mathscr{C}^k(K_n)$ are connected if and only if they are connected in K_n. So $\pi_n(p) = {}^k\tau^n_\pi(p)$ if $n \leq k$. This implies that for $n \leq k$ we can rewrite (3):

$$\begin{aligned} p^n_c(M) &= \sup\left\{p : {}^k\tau^n_\pi(p) < p^{site}_c\right\} \\ &\leq \sup\left\{p : {}^k\tau^\infty_\pi(p) < p^{site}_c\right\} = p_c(\Phi_{M,k}) \leq p_c(M), \end{aligned}$$

where we used that ${}^k\tau^n_\pi(p)$ decreases in n by Lemma 3. Theorem 2 states that $p^n_c(M)$ converges to $p_c(M)$, so we conclude that $\lim_{k\to\infty} p_c(\Phi_{M,k}) = p_c(M)$. ☐

5 Numerical results

In this section we present our numerical results. Our principal goal was to compute bounds using the alphabets $\mathscr{A}_{M,k}$ which were constructed in the previous section. In these calculations we encountered the problem of accumulating rounding errors, disturbing the convergence. Therefore we had to normalize the probability vector in each step. We did the rounding and normalization in such a way that our conclusions are not violated. For example to show that $\tau^n_\pi(p) < p^{site}_c$, we made sure that we computed an upper estimate for $\tau^n_\pi(p)$. Our implementation in Matlab (everything available from the author on request) gives the following results:

Proposition 2 *Take the alphabet $\mathscr{A}_{M,k}$ and define a classification $\mathscr{C}$ by the word code $\Phi_{M,k}$. Let $n = 1000$ and define $\tau^n_\pi(p)$ as before. Then*

- *For $M = 2$ and $k = 0$, we have $\tau^n_\pi(0.785) < p^{site}_c$.*
- *For $M = 2$ and $k = 1$, we have $\tau^n_\pi(0.859) < p^{site}_c$.*
- *For $M = 3$ and $k = 0$, we have $\tau^n_\pi(0.715) < p^{site}_c$.*

Corollary 2 $p_c(2) > 0.859$ *and* $p_c(3) > 0.715$.



Figure 5: Plot of $\tau_\pi^n(p)$ for $p = 0.7 + 0.01k$ where $k = 0, \dots, 9$ as functions of n. Note that there is a kind of phase transition between $p = 0.78$ and $p = 0.79$.

Proof This follows from Lemma 4 and Theorem 1. □

Figure 5 illustrates for the case $M = 2$ and $k = 0$ how $\tau_\pi^n(p)$ behaves as a function of n for some values of p. The values of $\tau_\pi^n(p)$ were calculated by iterating the recursion of Lemma 2.

For larger values of k the computations were too complicated to perform in a reasonable computation time. For example, the alphabet $\mathscr{A}_{2,2}$ already contains 35357670 letters. Nevertheless we will show that it is possible to improve the bounds of Corollary 2 by taking other alphabets or word codes.

For $M = 2$, define the element set E by dividing the left and right side of the unit square into 4 equal pieces and the bottom and top side into 2 equal pieces. This leads to an alphabet $\mathscr{A}_E$ that is in some sense in between $\mathscr{A}_{2,1}$ and $\mathscr{A}_{2,2}$. Analogous to our previous approach, we define a classification by choosing a word code based on weak connectivity. For this classification we find $\tau_\pi^{50}(0.876) < p_c^{site}$, which implies $p_c(2) > 0.876$.

One can improve this even a bit more by taking the alphabet $\mathscr{A}_{2,2}$ and defining a word code $\tilde{\Phi}_{2,2}$ that is a bit simpler than $\Phi_{2,2}$ as follows. If at least one of the letters in a 2×2 word w equals $\min(\mathscr{A}_{2,2})$, then $\tilde{\Phi}_{2,2}(w) = \Phi_{2,2}(w)$. Otherwise, define $\tilde{\Phi}_{2,2}(w)$ by first mapping each of the four letters to $\mathscr{A}_E$ and then mapping the new word to $\mathscr{A}_{2,2}$, in both steps using weak connectivity. This simplifies the required calculations a lot, and leads to $p_c(2) > 0.881$, in 200 iterations.

For $M = 3$ we improved the lower bound by using an alphabet that divides the left and right side of the unit square into 3 equal pieces. The bottom and top side still consist of one element. This leads to $p_c(3) > 0.784$, using 100 iterations.
These calculations have been checked by Arthur Bik, a mathematics student at Delft University of Technology. He independently implemented the algorithms and reproduced all results, except the bound $p_c(2) > 0.881$. This was due to the fact that his program was not fast enough to perform the calculations in a reasonable time. Concluding, the best lower bounds we found are

Theorem 3 *$p_c(2) > 0.881$ and $p_c(3) > 0.784$.*

Now let us turn to the upper bounds. Here we want to make use of Corollary 1. We already know $\tau^n_{\max(\mathscr{A})}(p) \leq \theta_n(p)$ for all n (Lemma 4) and that the classification is increasing. That means, we only have to find a probability vector $x \in [0,1]^{|\mathscr{A}|}$ for which $x_{\max(\mathscr{A})} > 0$ and $p \in (0,1]$ such that $F_{\mathscr{C}}(px + (1-p)\tau^{\square}) \succeq x$. The trick we use here is to take a value of p, and to search for the fixed point τ^{∞} by iterating the recursion of Lemma 2. Assume that the numerical results suggest that $\tau^{\infty}_{\max(\mathscr{A})} > 0$. Letting $x = \tau^n(p)$ for n large, we have a combination of x and p for which $F_{\mathscr{C}}(px + (1-p)\tau^{\square}) \approx x$. Taking p slightly larger gives a combination for which the desired inequality $F_{\mathscr{C}}(px+(1-p)\tau^{\square}) \succeq x$ holds. This strategy leads to the following results:

Proposition 3 *Take the alphabet $\mathscr{A}_{M,k}$ and define a classification $\mathscr{C}$ by the word code $\Psi_{M,k}$. Let $n = 1000$. The conditions $F_{\mathscr{C}}(px+(1-p)\tau^{\square}) \succeq x$ and $x_{\max(\mathscr{A})} > 0$ hold if x and p are chosen as follows:*

- *For $M = 3$ and $k = 0$, choose $p = 0.958$ and $x = \tau^n(0.9579)$.*
- *For $M = 4$ and $k = 0$, choose $p = 0.972$ and $x = \tau^n(0.9719)$.*

Corollary 3 *$p_c(3) < 0.958$ and $p_c(4) < 0.972$.*

Proof This follows from Lemma 4 and Corollary 1. □

For $M = 3$, the result can be sharpened by using the same alphabet that was used to improve the lower bound. The classification is again defined by a word code based on strong connectivity. In that case the choice $p = 0.940$ and $x = \tau^{1000}(0.9399)$ satisfies all conditions, so $p_c(3) < 0.940$.
The algorithm for $M = 4$ can be slightly adapted to find a bound for $M = 2$. Each realization of K_n for $M = 4$ can be seen as a realization of K_{2n} for $M = 2$. So the only thing that changes is the iteration function $F_{\mathscr{C}}$. Given a probability vector on $\mathscr{A}$, the probability on each 4×4 word can be calculated. The word code is still the same function $\Psi_{4,0}$. These ingredients

are sufficient to determine $F_{\mathscr{C}}$. Performing the calculations we find that the conditions are satisfied for $p = 0.993$ and $x = \tau^{1000}(0.9929)$, henceforth $p_c(2) < 0.993$.

Also for the upper bounds Arthur Bik checked our results. He independently reproduced our bounds, except for the bound $p_c(3) < 0.940$ (for similar reasons as before). Summarizing, our best upper bounds are

Theorem 4 *$p_c(2) < 0.993$, $p_c(3) < 0.940$ and $p_c(4) < 0.972$.*

Acknowledgement The author thanks Michel Dekking for the entertaining discussions on this topic and for his useful remarks on this paper. I am also grateful to Arthur Bik, who has spent a lot of time checking the numerical results. Figure 3 was made by Tilman Piesk [1].

References

[1] Berg, J. van den and Ermakov, A. – *A New Lower Bound for the Critical Probability of Site Percolation on the Square Lattice*, Random Structures and Algorithms, Vol. 8, No. 3 (1996).

[2] Chayes, J.T., Chayes, L. and Durrett, R. – *Connectivity Properties of Mandelbrot's Percolation Process*, Probability Theory and Related Fields, 77 (1988), 307–324.

[3] Dekking, F.M. and Grimmett, G.R. – *Superbranching processes and projections of random Cantor sets*, Probability Theory and Related Fields, 78 (1988), 335–355.

[4] Dekking, F.M. and Meester, R.W.J. – *On the Structure of Mandelbrot's Percolation and Other Random Cantor Sets*, Journal of Statistical Physics, Vol. 58, Nos. 5/6 (1990), 1109–1126.

[5] Falconer, K. – *Random fractals*, Math. Proc. Cambr. Phil. Soc. 100 (1986), 559-582.

[6] Grimmett G.R. – *Percolation*, Second edition, Springer-Verlag, Berlin (1999).

[7] Mandelbrot, B.B. – *The Fractal Geometry of Nature*, Freeman, San Francisco, 1983.

[8] Orzechowski, M. – *Geometrical and topological properties of fractal percolation*, PhD Thesis, University of St. Andrews, 1997.

[9] Wal, P. van der – *Random substitutions and fractal percolation*, PhD thesis, Delft University of Technology, 2002.

---

[1] License Terms: creativecommons.org/licenses/by/3.0/deed.en

[10] White, D.G. – *On the Value of the Critical Point in Fractal Percolation*, Random Structures and Algorithms, 18 (2001), 332–345.

Adress: TU Delft, EWI (DIAM), Section Applied Probability, Mekelweg 4,
2628 CD Delft, the Netherlands
e-mail: henkdon@gmail.com